

ON CLASSIFICATION OF COMPLEX FILIFORM LEIBNIZ ALGEBRAS.

J.R. GÓMEZ AND B.A. OMIROV

ABSTRACT. This work deal with Leibniz algebras. It is shown that in classifying of filiform non-Lie Leibniz algebras over the field of complex numbers which are obtained from the naturally graded non-Lie Leibniz algebras it suffices to consider some special basis transformations. Using this result, we derive a criterion that ascertains whether given two such filiform non-Lie Leibniz algebras are isomorphic in terms of such transformations.

2000 MSC: 17A32, 17B30.

Key-Words: *filiform Lie and Leibniz algebras, natural gradation, nilpotence, adapted basis.*

1. INTRODUCTION

The theory of Lie algebras is one of the most developed branches of modern algebra. It has been deeply investigated for many years by mathematicians, namely algebraists and specialists in representation theory, and by theoretical physicists. Active research of the properties of the Lie algebras leads to the investigation of the new and more general object - Leibniz algebras.

The present work is dedicated to the study of an algebraic object - Leibniz algebras, introduced into science by French mathematician J.-L. Loday [9] - [10] and investigated later in works [2] - [6], [11] - [12] and many others.

When investigating properties of the cyclic homologies, Loday had noted that if one replaces the external product in the definition of the n -th cochain by a tensor product, then in order to prove the differential property, that is defined on cochains, it suffices to show the validity of the Leibniz equality instead of the anticommutativity and the Jacoby equality. This was the motivation for the introduction of the Leibniz algebras, that are "noncommutative" generalization of the Lie algebras.

When studying a certain class of algebras it is important to describe at least the algebras of lower dimensions with the up to an isomorphism precision. For Leibniz algebras difficulties arise even when considering nilpotent algebras of dimension 5. Thus in this work we consider a special class of nilpotent Leibniz algebras, namely filiform algebras, that in the case of Lie algebras was studied in the works [7] - [8], [13]. In particular, in this work we devise a method of simplification of the basis transformations of the arbitrary filiform Leibniz algebras which were obtained from the naturally graded filiform non-Lie Leibniz algebras, that allows for consideration of only those transformations that change the structural constants in the description of such filiform Leibniz algebras [1]. An analogous method of simplification of the basis transformations in case of filiform Lie algebras was considered in the monograph [8]. The investigation of the non-Lie Leibniz filiform algebras which

were obtained from the naturally graded Lie algebra is the subject of the next paper.

2. PRELIMINARIES

Let L be an arbitrary Leibniz algebra of dimension n and $\{e_1, \dots, e_n\}$ be the basis of algebra L , then algebra L is defined by the products of basis elements, precisely $[e_i, e_j] = \sum_{k=1}^n \gamma_{ij}^k e_k$, where γ_{ij}^k are the structural constants. Thus the problem of classification becomes a problem of finding a description of structural constants up to a non-degenerate basis transformation.

From Leibniz equality we have the polynomial equalities:

$$\sum_{l=1}^n (\gamma_{jk}^l \gamma_{il}^m - \gamma_{ij}^l \gamma_{lk}^m + \gamma_{ik}^l \gamma_{lj}^m) = 0,$$

that define constraints set on the structural constants.

When solving classification problem for filiform Lie algebras in [8], it was shown that it is sufficient to consider only special types of transformations.

Recall that an algebra L over a field F is called a Leibniz algebra if it satisfies the following Leibniz identity:

$$[x, [y, z]] = [[x, y], z] - [[x, z], y],$$

where $[,]$ denotes the multiplication in L .

Note that if in the Leibniz algebra $[x, x] = 0$, the equality is true; then Leibniz equality is simplified into the Jacoby equality. Thus, the variety of Leibniz algebras is the "noncommutative" generalization of the Lie algebras' variety.

For a given arbitrary Leibniz algebra L we define the lower central series:

$$C^0(L) = L, \quad C^{k+1}(L) = [C^k(L), L].$$

Definition 2.1. A Leibniz algebra L is said to be filiform if $\dim C^i(L) = n - i - 1$, where $n = \dim L$ and $1 \leq i \leq n - 1$.

Since analogous results are known for Lie algebras [8], in the sequel we are going to consider non-Lie Leibniz algebras.

In [1] we obtained the decomposition of complex filiform Leibniz algebras which were obtained from naturally graded filiform non-Lie Leibniz algebras into two disjoint classes. Take into account the existence of filiform non-Lie Leibniz algebras which were obtained from the naturally graded Lie algebra; we have the following theorem:

Theorem 2.2. An arbitrary complex non-Lie filiform Leibniz algebra L of dimension $n + 1$ is isomorphic to one of the following algebras:

$$\mu_1^{\bar{\alpha}, \theta} = \begin{cases} [e_0, e_0] = e_2, \\ [e_i, e_0] = e_{i+1}, & 1 \leq i \leq n-1 \\ [e_0, e_1] = \alpha_3 e_3 + \alpha_4 e_4 + \dots + \alpha_{n-1} e_{n-1} + \theta e_n, \\ [e_j, e_1] = \alpha_3 e_{j+2} + \alpha_4 e_{j+3} + \dots + \alpha_{n+1-j} e_n, & 1 \leq j \leq n-2 \end{cases}$$

(omitted products are equal to zero)

$$\mu_2^{\bar{\beta}, \gamma} = \begin{cases} [e_0, e_0] = e_2, \\ [e_i, e_0] = e_{i+1}, & 2 \leq i \leq n-1 \\ [e_0, e_1] = \beta_3 e_3 + \beta_4 e_4 + \dots + \beta_n e_n, \\ [e_1, e_1] = \gamma e_n, \\ [e_j, e_1] = \beta_3 e_{j+2} + \beta_4 e_{j+3} + \dots + \beta_{n+1-j} e_n, & 2 \leq j \leq n-2 \end{cases}$$

(omitted products are equal to zero)

$$\mu_3^{\alpha, \beta, \delta} = \begin{cases} [e_0, e_0] = e_n, \\ [e_1, e_1] = \alpha e_n, \\ [e_i, e_0] = e_{i+1}, & 1 \leq i \leq n-1 \\ [e_0, e_1] = -e_2 + \beta e_n, \\ [e_0, e_i] = -e_{i+1}, & 2 \leq i \leq n-1 \\ [e_i, e_j] = -[e_j, e_i] \in \text{lin} < e_{i+j+1}, e_{i+j+2}, \dots, e_n >, & 1 \leq i \leq n-3, \\ & 2 \leq j \leq n-1-i \\ [e_{n-i}, e_i] = -[e_i, e_{n-i}] = (-1)^i \delta e_n, & 1 \leq i \leq n-1 \end{cases}$$

(omitted products are equal to zero)

where $[\cdot, \cdot]$ is the multiplication in L and $\{e_0, e_1, e_2, \dots, e_n\}$ is the basis of the algebra, $\delta \in \{0, 1\}$ for odd n and $\delta = 0$ for even n .

Remark 2.3. In the description of theorem 2.2 the isomorphisms inside the classes are unknown.

3. ON TRANSFORMATIONS OF COMPLEX FILIFORM LEIBNIZ ALGEBRAS.

Since an arbitrary non-Lie filiform Leibniz algebra up to an isomorphism belongs to one of the classes of the theorem 2.2, we conclude that to consider the isomorphisms inside the classes we need to study the behavior of the parameters under the action of the non-degenerate change of basis. As we mentioned above we will consider only the first two classes of the theorem 2.2.

Definition 3.1. Let L be a complex filiform $(n+1)$ -dimensional Leibniz algebra which is obtained from the naturally graded filiform non-Lie Leibniz algebras. Basis $\{e_0, e_1, \dots, e_n\}$ of algebra L is said to be adapted if the multiplication has the form $\mu_1^{\bar{\alpha}, \theta}$ or $\mu_2^{\bar{\beta}, \gamma}$.

Let L be the Leibniz algebra defined on a vector space V and $\{e_0, e_1, \dots, e_n\}$ is the adapted basis of the algebra L .

Definition 3.2. The transformation of the basis $f \in GL(V)$ is said to be adapted for the product of the algebra L if the basis $\{f(e_0), f(e_1), \dots, f(e_n)\}$ is adapted.

The closed subgroup of group $GL(V)$ spanned by adapted transformations will be denoted by $GL_{ad}(V)$.

In the sequel we will need the following lemma:

Lemma 3.3. For any $0 \leq p \leq n-k, 3 \leq k \leq n$ the following equality holds:

$$\sum_{i=k}^n a(i) \sum_{j=i+p}^n b(i, j) e_j = \sum_{j=k+p}^n \sum_{i=k}^{j-p} a(i) b(i, j) e_j \quad (1)$$

Proof. The following chain of equalities:

$$\begin{aligned}
& \sum_{i=k}^n a(i) \sum_{j=i+p}^n b(i, j) e_j = \\
&= \sum_{i=k}^n a(i) (b(i, i+p) e_{i+p} + b(i, i+p+1) e_{i+p+1} + \dots + b(i, n) e_n) = \\
&= a(k) (b(k, k+p) e_{k+p} + b(k, k+p+1) e_{k+p+1} + \dots + b(k, n) e_n) + \dots + \\
&+ a(t) (b(t, t+p) e_{t+p} + b(t, t+p+1) e_{t+p+1} + \dots + b(t, n) e_n) + \dots + \\
&+ a(n-p) b(n-p, n) e_n = a(k) b(k, k+p) e_{k+p} + (a(k) b(k, k+p+1) + \\
&+ a(k+1) b(k+1, k+p+1)) e_{k+p+1} + \dots + (a(k) b(k, t+p) + \\
&+ a(k+1) b(k+1, t+p) + \dots + a(t) b(t, t+p)) e_{t+p} + \dots + \\
&+ a(n-p) b(n-p, n) e_n = \sum_{t=k}^{n-p} \sum_{i=k}^t a(i) b(i, t+p) e_{t+p} = \sum_{j=k+p}^n \sum_{i=k}^{j-p} a(i) b(i, j) e_j
\end{aligned}$$

proves the lemma. $\square$

Proposition 3.4. *Let $f \in GL_{ad}(V)$.*

a) *If the algebra L belongs to the class $\mu_1^{\bar{\alpha}, \theta}$ in theorem 2.2, then f has the following form:*

$$\begin{cases} f(e_0) = a_0 e_0 + a_1 e_1 + a_2 e_2 + \dots + a_n e_n \\ f(e_1) = (a_0 + a_1) e_1 + a_2 e_2 + \dots + a_{n-2} e_{n-2} + (a_{n-1} + a_1(\theta - \alpha_n)) e_{n-1} + b_n e_n \\ f(e_{i+1}) = [f(e_i), f(e_0)], \quad 1 \leq i \leq n-1 \\ f(e_2) = [f(e_0), f(e_0)]. \end{cases}$$

b) *If the algebra L belongs to the class $\mu_2^{\bar{\beta}, \gamma}$ in theorem 2.2, then f has the following form:*

$$\begin{cases} f(e_0) = a_0 e_0 + a_1 e_1 + a_2 e_2 + \dots + a_n e_n \\ f(e_1) = b_1 e_1 - \frac{a_1 b_1 \gamma}{a_0} e_{n-1} + b_n e_n \\ f(e_{i+1}) = [f(e_i), f(e_0)], \quad 2 \leq i \leq n-1 \\ f(e_2) = [f(e_0), f(e_0)]. \end{cases}$$

Proof. Let $f \in GL_{ad}(V)$.

Set $f(e_0) = a_0 e_0 + a_1 e_1 + \dots + a_n e_n$, $f(e_1) = b_0 e_0 + b_1 e_1 + \dots + b_n e_n$.

We prove case a).

Consider the product: $f(e_2) = [f(e_0), f(e_0)]$.

Using (1) we have the equalities:

$$\begin{aligned}
& [f(e_0), f(e_0)] = [a_0 e_0 + a_1 e_1 + \dots + a_n e_n, a_0 e_0 + a_1 e_1 + \dots + a_n e_n] = a_0(a_0 + a_1) e_2 + \\
& + a_0 \sum_{i=3}^n a_{i-1} e_i + a_0 a_1 \left(\sum_{i=3}^{n-1} \alpha_i e_i + \theta e_n \right) + a_1^2 \sum_{i=3}^n \alpha_i e_i + a_1 \sum_{i=2}^{n-2} a_i \sum_{k=i+2}^n \alpha_{k+1-i} e_k = \\
& = a_0(a_0 + a_1) e_2 + a_0 \sum_{i=3}^n a_{i-1} e_i + a_1(a_0 + a_1) \sum_{i=3}^{n-1} \alpha_i e_i + a_1(a_0 \theta + a_1 \alpha_n) e_n + \\
& + a_1 \sum_{i=4}^n a_{i-2} \sum_{k=i}^n \alpha_{k+3-i} e_k = a_0(a_0 + a_1) e_2 + a_0 \sum_{i=3}^n a_{i-1} e_i + a_1(a_0 + a_1) \sum_{i=3}^{n-1} \alpha_i e_i + \\
& + a_1(a_0 \theta + a_1 \alpha_n) e_n + a_1 \sum_{i=4}^n \sum_{k=i}^k (a_{i-2} \alpha_{k+3-i} e_k) = a_0(a_0 + a_1) e_2 + (a_0 a_2 + \\
& + a_1(a_0 + a_1) \alpha_3) e_3 + \sum_{t=4}^{n-1} (a_0 a_{t-1} + a_1(a_0 + a_1) \alpha_t + a_1 \sum_{i=4}^t a_{i-2} \alpha_{t+3-i}) e_t + \\
& + (a_0 a_{n-1} + a_1(a_0 \theta + a_1 \alpha_n) + a_1 \sum_{i=4}^n a_{i-2} \alpha_{n+3-i}) e_n.
\end{aligned}$$

Since $[f(e_0), f(e_0)] \in C^1(L)$, we have that $a_0(a_0 + a_1) \neq 0$, i.e. $a_0 \neq 0$ and $a_0 + a_1 \neq 0$.

Consider the product:

$$[f(e_0), f(e_1)] = b_0(a_0 + a_1)e_2 + \sum_{i=3}^n c_i e_i.$$

Since $[f(e_0), f(e_1)] \notin C^1(L)$ and $a_0 + a_1 \neq 0$, we have that $b_0 = 0$.

From the properties of adapted transformation we have that $f(e_2) = [f(e_1), f(e_0)]$.

Decomposition of the expression $[f(e_1), f(e_0)]$ in basis $e_0, e_1, \dots, e_n$ has the form:

$$\begin{aligned} [f(e_1), f(e_0)] &= a_0 b_1 e_2 + (a_0 b_2 + a_1 b_1 \alpha_3) e_3 + \sum_{t=4}^{n-1} (a_0 b_{t-1} + a_1 b_1 \alpha_t + a_1 \sum_{i=4}^t b_{i-2} \alpha_{t+3-i}) e_t + \\ &+ (a_0 b_{n-1} + a_1 b_1 \alpha_n + a_1 \sum_{i=4}^n b_{i-2} \alpha_{n+3-i}) e_n. \end{aligned}$$

Comparing the coefficients of the basis elements we get the following constraints of the coefficients of the transformation f :

$$\begin{cases} a_0 + a_1 = b_1 \\ a_2 = b_2 \\ a_0 a_{t-1} + a_1 \sum_{i=4}^t a_{i-2} \alpha_{t+3-i} = a_0 b_{t-1} + a_1 \sum_{i=4}^t b_{i-2} \alpha_{t+3-i} \\ a_0 a_{n-1} + a_1 (a_0 \theta + a_1 \alpha_n) + a_1 \sum_{i=4}^n a_{i-2} \alpha_{n+3-i} = a_0 b_{n-1} + a_1 b_1 \alpha_n + a_1 \sum_{i=4}^n b_{i-2} \alpha_{n+3-i}. \end{cases}$$

From these we have

$$\begin{cases} b_1 = a_0 + a_1 \\ b_i = a_i, \quad 2 \leq i \leq n-2 \\ b_{n-1} = a_{n-1} + a_1(\theta - \alpha_n). \end{cases}$$

Case b) is proved in the same way. $\square$

Analogously to [8], we introduce the notion of elementary transformations that are true for the non-Lie Leibniz algebras.

Definition 3.5. *The following types of basis transformations are said to be elementary:*

$$\text{first type} - \tau(a, b, k) = \begin{cases} f(e_0) = e_0 + a e_k \\ f(e_1) = e_1 + b e_k \\ f(e_{i+1}) = [f(e_i), f(e_0)], \quad 1 \leq i \leq n-1, \quad 2 \leq k \leq n \\ f(e_2) = [f(e_0), f(e_0)] \end{cases}$$

$$\text{second type} - \vartheta(a, b) = \begin{cases} f(e_0) = a e_0 + b e_1 \\ f(e_1) = (a + b) e_1 + b(\theta - \alpha_n) e_{n-1}, \quad a(a + b) \neq 0 \\ f(e_{i+1}) = [f(e_i), f(e_0)], \quad 1 \leq i \leq n-1, \\ f(e_2) = [f(e_0), f(e_0)] \end{cases}$$

$$\text{third type} - \sigma(b, n) = \begin{cases} f(e_0) = e_0 \\ f(e_1) = e_1 + b e_n, \\ f(e_{i+1}) = [f(e_i), f(e_0)], \quad 2 \leq i \leq n-1, \\ f(e_2) = [f(e_0), f(e_0)] \end{cases}$$

$$\text{fourth type} - \eta(a, k) = \begin{cases} f(e_0) = e_0 + a e_k \\ f(e_1) = e_1 \\ f(e_{i+1}) = [f(e_i), f(e_0)], \quad 2 \leq i \leq n-1, \quad 2 \leq k \leq n, \\ f(e_2) = [f(e_0), f(e_0)] \end{cases}$$

$$\text{fifth type} - \delta(a, b, d) = \begin{cases} f(e_0) = ae_0 + be_1 \\ f(e_1) = de_1 - \frac{bd\gamma}{a}e_{n-1}, & ad \neq 0 \\ f(e_{i+1}) = [f(e_i), f(e_0)], & 2 \leq i \leq n-1, \\ f(e_2) = [f(e_0), f(e_0)] \end{cases}$$

where $a, b, d \in C$.

Let f be an arbitrary element of the group $GL_{ad}(V)$, then f is expressed via the elementary transformations, namely the following is true:

Proposition 3.6.

i) Let f have the form from the proposition 3.4 a). Then

$$f = \tau(a_n, b_n, n) \circ \tau(a_{n-1}, a_{n-1}, n-1) \circ \dots \circ \tau(a_2, a_2, 2) \circ \vartheta(a_0, a_1)$$

ii) Let f have the form from proposition 3.4 b). Then

$$f = \sigma(b_n, n) \circ \eta(a_n, n) \circ \eta(a_{n-1}, n-2) \circ \dots \circ \eta(a_2, 2) \circ \delta(a_0, a_1, b_1)$$

Proof. The proposition is validated straightforwardly. $\square$

For the aforementioned decompositions the following is true:

Proposition 3.7.

1) Transformation $g = \tau(a_n, b_n, n) \circ \tau(a_{n-1}, a_{n-1}, n-1) \circ \dots \circ \tau(a_2, a_2, 2)$ does not change the structural constants of the first class of algebras of the theorem 2.2.

2) Transformation $\varphi = \sigma(b_n, n) \circ \eta(a_n, n) \circ \eta(a_{n-1}, n-2) \circ \dots \circ \eta(a_2, 2)$ does not change the structural constants of the second class of algebras from the theorem 2.2.

Proof. Let us prove the first assertion.

Consider the transformation $\tau(a, b, k)$:

$$\tau(a, b, k) = \begin{cases} f(e_0) = e_0 + ae_k \\ f(e_1) = e_1 + be_k \\ f(e_{i+1}) = [f(e_i), f(e_0)], & 1 \leq i \leq n-1, \quad 2 \leq k \leq n \\ f(e_2) = [f(e_0), f(e_0)] \end{cases}$$

When $2 \leq k \leq n-1$ set $a = b$ then $\tau(a, b, k)$ takes the following form:

$$\tau(a, a, k) = \begin{cases} f(e_0) = e_0 + ae_k \\ f(e_1) = e_1 + ae_k \\ f(e_{i+1}) = [f(e_i), f(e_0)], & 1 \leq i \leq n-1, \\ f(e_2) = [f(e_0), f(e_0)] \end{cases}$$

Note that this transformation is adapted. In fact, when considering the products:

$$\begin{aligned} f(e_2) &= [f(e_0), f(e_0)] = [e_0 + ae_k, e_0 + ae_k] = e_2 + ae_{k+1}, \\ f(e_2) &= [f(e_1), f(e_0)] = [e_1 + ae_k, e_0 + ae_k] = e_2 + ae_{k+1}, \\ f(e_i) &= [f(e_{i-1}), f(e_0)] = [e_{i-1} + ae_{k+i-2}, e_0 + ae_k] = e_i + ae_{k+1}, \text{ when } k+i-2 \leq n, \\ &\text{otherwise we suppose } a = 0 \text{ and thus we find that } \tau(a, a, k) \in GL_{ad}(V). \end{aligned}$$

Consider the products in which the parameters are defined:

$$\begin{aligned} [f(e_0), f(e_1)] &= [e_0 + ae_k, e_1 + ae_k] = \sum_{i=3}^{n-1} \alpha_i e_i + \theta e_n + a \sum_{i=3}^{n-k+1} \alpha_i e_{k+i-1} = \\ &= \sum_{i=3}^{n-k+1} \alpha_i (e_i + ae_{k+i-1}) + \sum_{i=n-k+2}^{n-1} \alpha_i e_i + \theta e_n = \sum_{i=3}^{n-1} \alpha_i f(e_i) + \theta f(e_n), \\ [f(e_1), f(e_1)] &= [e_1 + ae_k, e_1 + ae_k] = \sum_{i=3}^n \alpha_i e_i + a \sum_{i=3}^{n-k+1} \alpha_i e_{k+i-1} = \\ &= \sum_{i=3}^{n-k+1} \alpha_i (e_i + ae_{k+i-1}) + \sum_{i=n-k+2}^n \alpha_i e_i = \sum_{i=3}^n \alpha_i f(e_i). \end{aligned}$$

Thus the transformations of the first type for $2 \leq k \leq n-1$ do not change the parameters α_i, θ for arbitrary $a \in C$.

Analogously, we check that $\tau(a, b, n) \in GL_{ad}(V)$ does not change parameters α_i, θ for arbitrary $a \in C$.

Since superposition of adapted transformations is again an adapted transformation, we can conclude that transformation $g = \tau(a_n, b_n, n) \circ \tau(a_{n-1}, a_{n-1}, n-1) \circ \dots \circ \tau(a_2, a_2, 2)$ does not change the structural constants of class $\mu_1^{\bar{\alpha}\theta}$ from the theorem 2.2.

The proof of the second assertion of the proposition is carried out in similar way. $\square$

Thus, we derived that the problem of the classification is reduced to the problem of investigation of elementary transformations of second and fifth types for classes $\mu_1^{\bar{\alpha},\theta}$ and $\mu_2^{\bar{\beta},\gamma}$, respectively.

4. CRITERION OF ISOMORPHISMS OF COMPLEX FILIFORM NON-LIE LEIBNIZ ALGEBRAS.

For an arbitrary element a of the Leibniz algebra L , denote the operator of right multiplication by $R_a(x)$, e.g. $R_a(x) = [x, a]$ for any $x \in L$.

Set $R_a^m(x) := [\underbrace{\dots [x, a], a], \dots, a]}_{m\text{-times}}$ and $R_a^0(x) := x$.

It is easy to see that in algebra L , if it has the form of the first two classes of theorem 2.2, we have the equality:

$$[[e_s, e_1], e_0] = [e_{s+1}, e_1], \quad (2 \leq s \leq n). \quad (2)$$

Let L be the algebra that belongs to class $\mu_1^{\bar{\alpha},\theta}$ (class $\mu_2^{\bar{\beta},\gamma}$). Then from (2) we have that for arbitrary $m \in N$, $0 \leq p \leq n$ (respectively $0 \leq p \leq n$, $p \neq 1$) the following equality holds true:

$$R_{e_1}^m(e_p) = R_{e_0}^{p-1}(R_{e_1}^m(e_0)). \quad (3)$$

To prove the main theorem of this paragraph we will need the following results:

Lemma 4.1. *Let L be a non-Lie filiform Leibniz algebra of the first two classes from the theorem 2.2. Then when $2 \leq m \leq \frac{n-1}{2}$ the following equality is true:*

$$R_{e_1}^m(e_0) = \sum_{i_m=2m+1}^n \sum_{i_{m-1}=2m+1}^{i_m} \dots \sum_{i_1=2m+1}^{i_2} \eta_{i_m+3-i_{m-1}} \cdot \dots \cdot \eta_{i_2+3-i_1} \cdot \eta_{i_1+3-2(m-1)} e_{i_m},$$

$$\text{where } \eta_i = \begin{cases} \alpha_i, & \text{when } L \text{ belongs to the first class} \\ \beta_i, & \text{when } L \text{ belongs to the second class.} \end{cases}$$

Proof. Let $\eta_i = \alpha_i$. The case when $\eta_i = \beta_i$ is similar.

We will carry the proof using the method of mathematical induction and (1).

When $m = 2$, we have

$$\begin{aligned} R_{e_1}^2(e_0) &= \left[\sum_{i=3}^{n-1} \alpha_i e_i + \theta e_n, e_1 \right] = \sum_{i=5}^n \alpha_{i-2} [e_{i-2}, e_1] = \sum_{i=5}^n \alpha_{i-2} \sum_{j=i}^n \alpha_{j+3-i} e_1 = \\ &= \sum_{j=5}^n \sum_{i=5}^j \alpha_{i-2} \alpha_{j+3-i} e_1. \end{aligned}$$

Assume that the equality is true for $m = k$, e.g. the following equality:

$$R_{e_1}^k(e_0) = \sum_{i_k=2k+1}^n \sum_{i_{k-1}=2k+1}^{i_k} \dots \sum_{i_1=2k+1}^{i_2} \alpha_{i_k+3-i_{k-1}} \cdot \dots \cdot \alpha_{i_2+3-i_1} \cdot \alpha_{i_1+3-2(k-1)} e_{i_k}.$$

Let us proof the equality for $m = k + 1$. Using the induction hypothesis and the chain of equalities:

$$\begin{aligned}
R_{e_1}^{k+1}(e_0) &= [R_{e_1}^k(e_0), e_1] = \\
&= \left[\sum_{i_k=2k+1}^n \sum_{i_{k-1}=2k+1}^{i_k} \dots \sum_{i_1=2k+1}^{i_2} \alpha_{i_k+3-i_{k-1}} \cdot \dots \cdot \alpha_{i_2+3-i_1} \cdot \alpha_{i_1+3-(2k+1)} e_{i_k}, e_1 \right] = \\
&= \left[\sum_{i_k=2k+3}^n \sum_{i_{k-1}=2k+3}^{i_k} \dots \sum_{i_1=2k+3}^{i_2} \alpha_{i_k+3-i_{k-1}} \cdot \dots \cdot \alpha_{i_2+3-i_1} \cdot \alpha_{i_1+3-(2k+3)} e_{i_k-2}, e_1 \right] = \\
&= \sum_{i_k=2k+3}^n \sum_{i_{k-1}=2k+3}^{i_k} \dots \sum_{i_1=2k+3}^{i_2} \alpha_{i_k+3-i_{k-1}} \cdot \dots \cdot \alpha_{i_2+3-i_1} \cdot \\
&\quad \cdot \alpha_{i_1+3-(2k+3)} \sum_{i_{k+1}=i_k}^n \alpha_{i_{k+1}+3-i_k} e_{i_{k+1}} = \\
&= \sum_{i_k=2k+3}^n \sum_{i_{k+1}=i_k}^n \sum_{i_{k-1}=2k+3}^{i_k} \dots \sum_{i_1=2k+3}^{i_2} \alpha_{i_k+3-i_{k-1}} \cdot \dots \cdot \alpha_{i_2+3-i_1} \cdot \alpha_{i_1+3-(2k+3)} e_{i_{k+1}} = \\
&= \sum_{i_{k+1}=2k+3}^n \sum_{i_k=2k+3}^{i_{k+1}} \sum_{i_{k-1}=2k+3}^{i_k} \dots \sum_{i_1=2k+3}^{i_2} \alpha_{i_k+3-i_{k-1}} \cdot \dots \cdot \alpha_{i_2+3-i_1} \cdot \alpha_{i_1+3-(2k+3)} e_{i_{k+1}}
\end{aligned}$$

we obtain the induction proof. $\square$

From the first assertion of the proposition 3.7 we have that the adapted transformations for the class $\mu_1^{\alpha, \theta}$ are reduced to the transformations of the form:

$$\begin{cases} e'_0 = Ae_0 + Be_1 \\ e'_1 = (A+B)e_1 + B(\theta - \alpha_n)e_{n-1}, \end{cases} \quad (4)$$

where $A(A+B) \neq 0$.

Under the action of the given basis change we have

Corollary 4.2. For arbitrary $3 \leq k \leq n$,

$$e'_k = (A+B) \left(\sum_{i=0}^{k-2} C_{k-1}^{k-1-i} A^{k-1-i} B^i R_{e_1}^i(e_{k-i}) + B^{k-1} R_{e_1}^{k-1}(e_0) \right) \quad (5)$$

Proof.

We shall use the method of mathematical induction.

When $k = 3$ we have:

$$\begin{aligned}
e'_3 &= [e'_2, e'_0] = [A(A+B)e_2 + B(A+B) \sum_{i=3}^{n-1} \alpha_i e_i + B(A\theta + B\alpha_n), Ae_0 + Be_1] = \\
&= A^2(A+B)e_3 + 2AB(A+B)[e_2, e_1] + B^2(A+B)[[e_0, e_1], e_1] = (A+B)(A^2e_3 + \\
&\quad + 2ABR_{e_1}(e_2) + B^2R_{e_1}^2(e_0)), \text{ i.e. (4) is true.}
\end{aligned}$$

Suppose that the equality (5) is true for $k=p$, i.e.

$$e'_p = (A+B) \left(\sum_{i=0}^{p-2} C_{p-1}^{p-1-i} A^{p-1-i} B^i R_{e_1}^i(e_{p-i}) + B^{p-1} R_{e_1}^{p-1}(e_0) \right)$$

We shall now prove the equality (5) for $k = p + 1$. Taking into account equality (2) and the following chain of equalities:

$$\begin{aligned}
e'_{p+1} &= [e'_p, e_0] = \\
&= [(A+B) \left(\sum_{i=0}^{p-2} C_{p-1}^{p-1-i} A^{p-1-i} B^i R_{e_1}^i(e_{p-i}) + B^{p-1} R_{e_1}^{p-1}(e_0) \right), Ae_0 + Be_1] = \\
&= (A+B) \left(\sum_{i=0}^{p-2} C_{p-1}^{p-1-i} A^{p-i} B^i R_{e_1}^i(e_{p+1-i}) + AB^{p-1} R_{e_1}^{p-1}(e_2) + \right. \\
&\quad \left. + \sum_{i=0}^{p-2} C_{p-1}^{p-1-i} A^{p-1-i} B^{i+1} R_{e_1}^{i+1}(e_{p-i}) + B^p R_{e_1}^p(e_0) \right) =
\end{aligned}$$

$$\begin{aligned}
&= (A+B) \cdot \left(\sum_{i=0}^{p-2} C_{p-1}^{p-1-i} A^{p-i} B^i R_{e_1}^i(e_{p+1-i}) + AB^{p-1} R_{e_1}^{p-1}(e_2) + \right. \\
&+ \left. \sum_{i=1}^{p-1} C_{p-1}^{p-i} A^{p-i} B^i R_{e_1}^i(e_{p+1-i}) + B^p R_{e_1}^p(e_0) \right) = \\
&= (A+B) \left(\sum_{i=1}^{p-2} (C_{p-1}^{p-1-i} + C_{p-1}^{p-i}) A^{p-i} B^i R_{e_1}^i(e_{p+1-i}) + \right. \\
&+ \left. C_{p-1}^{p-1} A^p e_{p+1} + C_{p-1}^1 AB^{p-1} R_{e_1}^{p-1}(e_2) + AB^{p-1} R_{e_1}^{p-1}(e_2) + \right. \\
&+ \left. B^p R_{e_1}^p(e_0) \right) = (A+B) \left(\sum_{i=1}^{p-2} C_p^{p-i} A^{p-i} B^i R_{e_1}^i(e_{p+1-i}) + \right. \\
&+ \left. C_p^p A^p e_{p+1} + C_p^1 AB^{p-1} R_{e_1}^{p-1}(e_2) + B^p R_{e_1}^p(e_0) \right) = \\
&= (A+B) \left(\sum_{i=0}^{p-1} C_p^{p-i} A^{p-i} B^i R_{e_1}^i(e_{p+1-i}) + B^p R_{e_1}^p(e_0) \right)
\end{aligned}$$

we complete the proof. $\square$

From the second assertion of the proposition 3.7 we have that the adapted transformations for the class $\mu_2^{\bar{\beta}, \gamma}$ are reduced to transformations of the form:

$$\begin{cases} e'_0 = Ae_0 + Be_1 \\ e'_1 = De_1 - \frac{BD\gamma}{A} e_{n-1}, \end{cases}$$

where $AD \neq 0$.

Under the action of the above basis change following is true

Corollary 4.3. *For arbitrary $3 \leq k \leq n$*

$$e'_k = A \left(\sum_{i=0}^{k-2} C_{k-1}^{k-1-i} A^{k-1-i} B^i R_{e_1}^i(e_{k-i}) + B^{k-1} R_{e_1}^{k-1}(e_0) \right)$$

Proof. The proof is carried out in a similar way to the proof of the corollary 4.2. $\square$

We will write an algebra from class $\mu_1^{\bar{\alpha}, \theta}$ ($\mu_2^{\bar{\beta}, \gamma}$) in the form of $L(\alpha_3, \alpha_4, \dots, \alpha_n, \theta)$ (respectively $L(\beta_3, \beta_4, \dots, \beta_n, \gamma)$).

Theorem 4.4.

a) *Two algebras $L(\alpha_3, \alpha_4, \dots, \alpha_n, \theta)$ and $L'(\alpha'_3, \alpha'_4, \dots, \alpha'_n, \theta')$ are isomorphic if and only if there exist $A, B \in C$ such that $A(A+B) \neq 0$ and the following conditions hold:*

$$\begin{aligned}
\alpha'_3 &= \frac{(A+B)}{A^2} \alpha_3 \\
\alpha'_t &= \frac{1}{A^{t-1}} \left((A+B) \alpha_t - \sum_{k=3}^{t-1} (C_{k-1}^{k-2} A^{k-2} B \alpha_{t+2-k} + \right. \\
&+ C_{k-1}^{k-3} A^{k-3} B^2 \sum_{i_1=k+2}^t \alpha_{t+3-i_1} \alpha_{i_1+1-k} + \\
&+ C_{k-1}^{k-4} A^{k-4} B^3 \sum_{i_2=k+3}^t \sum_{i_1=k+3}^{i_2} \alpha_{t+3-i_2} \cdot \alpha_{i_2+3-i_1} \cdot \alpha_{i_1-k} + \dots + \\
&+ C_{k-1}^1 AB^{k-2} \sum_{i_{k-3}=2k-2}^t \sum_{i_{k-4}=2k-2}^{i_{k-3}} \dots \sum_{i_1=2k-2}^{i_2} \alpha_{t+3-i_{k-3}} \cdot \alpha_{i_{k-3}+3-i_{k-4}} \cdot \dots \cdot \alpha_{i_2+3-i_1} \alpha_{i_1+5-2k} + \\
&+ B^{k-1} \sum_{i_{k-2}=2k-1}^t \sum_{i_{k-3}=2k-1}^{i_{k-2}} \dots \sum_{i_1=2k-1}^{i_2} \alpha_{t+3-i_{k-2}} \cdot \alpha_{i_{k-2}+3-i_{k-3}} \cdot \dots \cdot \alpha_{i_2+3-i_1} \cdot \alpha_{i_1+4-2k} \Big) \cdot \alpha'_k \Big),
\end{aligned}$$

$$\begin{aligned} \theta' = & \frac{1}{A^{n-1}} \left(A\theta + B\alpha_n - \sum_{k=3}^{n-1} (C_{k-1}^{k-2} A^{k-2} B\alpha_{n+2-k} + C_{k-1}^{k-3} A^{k-3} B^2 \sum_{i_1=k+2}^n \alpha_{n+3-i_1} \alpha_{i_1+1-k} + \right. \\ & + C_{k-1}^{k-4} A^{k-4} B^3 \sum_{i_2=k+3}^n \sum_{i_1=k+3}^{i_2} \alpha_{n+3-i_2} \alpha_{i_2+3-i_1} \alpha_{i_1-k} + \dots + \\ & + C_{k-1}^1 A B^{k-2} \sum_{i_{k-3}=2k-2}^n \sum_{i_{k-4}=2k-2}^{i_{k-3}} \dots \sum_{i_1=2k-2}^{i_2} \alpha_{n+3-i_{k-3}} \alpha_{i_{k-3}+3-i_{k-4}} \dots \alpha_{i_2+3-i_1} \alpha_{i_1+5-2k} + \\ & \left. + B^{k-1} \sum_{i_{k-2}=2k-1}^n \sum_{i_{k-3}=2k-1}^{i_{k-2}} \dots \sum_{i_1=2k-1}^{i_2} \alpha_{n+3-i_{k-2}} \cdot \alpha_{i_{k-2}+3-i_{k-3}} \cdot \dots \cdot \alpha_{i_2+3-i_1} \cdot \alpha_{i_1+4-2k} \right) \cdot \alpha'_k \Big), \end{aligned}$$

where $4 \leq t \leq n$.

b) Two algebras $L(\beta_3, \beta_4, \dots, \beta_n, \gamma)$ and $L'(\beta'_3, \beta'_4, \dots, \beta'_n, \gamma')$ are isomorphic if and only if there exist $A, B \in C$ such that $A(A+B) \neq 0$ and the following conditions hold:

$$\begin{aligned} \gamma' &= \frac{D^2}{A^n} \gamma \\ \beta'_3 &= \frac{D}{A^2} \beta_3 \\ \beta'_t &= \frac{1}{A^{t-1}} (D\beta_t - \sum_{k=3}^{t-1} (C_{k-1}^{k-2} A^{k-2} B\beta_{t+2-k} + C_{k-1}^{k-3} A^{k-3} B^2 \sum_{i_1=k+2}^t \beta_{t+3-i_1} \cdot \beta_{i_1+1-k} + \\ & + C_{k-1}^{k-4} A^{k-4} B^3 \sum_{i_2=k+3}^t \sum_{i_1=k+3}^{i_2} \beta_{t+3-i_2} \cdot \beta_{i_2+3-i_1} \cdot \beta_{i_1-k} + \dots + \\ & + C_{k-1}^1 A B^{k-2} \sum_{i_{k-3}=2k-2}^t \sum_{i_{k-4}=2k-2}^{i_{k-3}} \dots \sum_{i_1=2k-2}^{i_2} \beta_{t+3-i_{k-3}} \beta_{i_{k-3}+3-i_{k-4}} \dots \beta_{i_2+3-i_1} \beta_{i_1+5-2k} + \\ & + B^{k-1} \sum_{i_{k-2}=2k-1}^t \sum_{i_{k-3}=2k-1}^{i_{k-2}} \dots \sum_{i_1=2k-1}^{i_2} \beta_{t+3-i_{k-2}} \beta_{i_{k-2}+3-i_{k-3}} \dots \beta_{i_2+3-i_1} \beta_{i_1+4-2k}) \beta'_k), \end{aligned}$$

where $4 \leq t \leq n-1$.

$$\begin{aligned} \beta'_n &= \frac{BD\gamma}{A^n} + \frac{1}{A^{n-1}} (D\beta_n - \sum_{k=3}^{n-1} (C_{k-1}^{k-2} A^{k-2} B\beta_{n+2-k} + C_{k-1}^{k-3} A^{k-3} B^2 \sum_{i_1=k+2}^n \beta_{n+3-i_1} \cdot \beta_{i_1+1-k} + \\ & + C_{k-1}^{k-4} A^{k-4} B^3 \sum_{i_2=k+3}^n \sum_{i_1=k+3}^{i_2} \beta_{n+3-i_2} \cdot \beta_{i_2+3-i_1} \cdot \beta_{i_1-k} + \dots + \\ & + C_{k-1}^1 A B^{k-2} \sum_{i_{k-3}=2k-2}^n \sum_{i_{k-4}=2k-2}^{i_{k-3}} \dots \sum_{i_1=2k-2}^{i_2} \beta_{n+3-i_{k-3}} \beta_{i_{k-3}+3-i_{k-4}} \dots \beta_{i_2+3-i_1} \beta_{i_1+5-2k} + \\ & + B^{k-1} \sum_{i_{k-2}=2k-1}^n \sum_{i_{k-3}=2k-1}^{i_{k-2}} \dots \sum_{i_1=2k-1}^{i_2} \beta_{n+3-i_{k-2}} \beta_{i_{k-2}+3-i_{k-3}} \dots \beta_{i_2+3-i_1} \beta_{i_1+4-2k}) \beta'_k), \end{aligned}$$

Remark 4.5. From theorem 4.4 we have that α'_k is a polynomial with the unknown coefficients A, B that satisfy to the condition $A(A+B) \neq 0$ and with the given $\alpha_1, \alpha_2, \dots, \alpha_k, \alpha'_1, \alpha'_2, \dots, \alpha'_{k-1}$. And β'_k is a polynomial with the unknown coefficients A, B, D that satisfy to the condition $AD \neq 0$ and with the given $\beta_1, \beta_2, \dots, \beta_k, \beta'_1, \beta'_2, \dots, \beta'_{k-1}$, i.e. finding parameters α'_k and β'_k is a recursive procedure.

Thus, we can conclude that in any dimension the problem of the classification of complex filiform Leibniz algebras which are obtained from the naturally graded filiform non-Lie Leibniz algebras up to an isomorphism is algorithmically solvable.

Proof of the theorem.

Consider the class $\mu_1^{\bar{\alpha}, \theta}$. Let $\{e_0, e_1, \dots, e_n\}$ be the basis of $L(\alpha_3, \alpha_4, \dots, \alpha_n, \theta)$, and $\{e'_0, e'_1, \dots, e'_n\}$ be the basis of $L'(\alpha'_3, \alpha'_4, \dots, \alpha'_n, \theta')$.

It is easy to see that in algebra $L(\alpha_3, \alpha_4, \dots, \alpha_n, \theta)$ the following is true:

$$[[e_0, e_1], e_1] = [[e_1, e_1], e_1].$$

We will consider the change of basis (4).

From lemma 4.1 and equality (3) we obtain

$$R_{e_1}^m(e_{k-m}) = \sum_{i_m=k+m}^n \sum_{i_{m-1}=k+m}^{i_m} \dots \sum_{i_1=2m+1}^{i_2} \alpha_{i_m+3-i_{m-1}} \dots \alpha_{i_2+3-i_1} \alpha_{i_1+3-(k+m)} e_{i_m} \quad (6)$$

where $m \leq n-k$ $m \leq k \leq n$.

Using (6) in equality (5) we can carry following changes:

$$\begin{aligned} e'_k &= (A+B) \left(\sum_{i=0}^{k-2} C_{k-1}^{k-1-i} A^{k-1-i} B^i R_{e_1}^i(e_{k-1}) + B^{k-1} R_{e_1}^{k-1}(e_0) \right) = (A+B)(A^{k-1}e_k + \\ &+ C_{k-1}^{k-2} A^{k-2} B \sum_{i=k+1}^n \alpha_{i+2-k} e_i + C_{k-1}^{k-3} A^{k-3} B^2 \sum_{i=k+2}^n \sum_{i_1=k+2}^{i_2} \alpha_{i+3-i_1} \cdot \alpha_{i_1+1-k} e_i + \\ &+ C_{k-1}^{k-4} A^{k-4} B^3 \sum_{i=k+3}^n \sum_{i_2=k+3}^{i_1} \sum_{i_1=k+3}^{i_2} \alpha_{i+3-i_2} \cdot \alpha_{i_2+3-i_1} \cdot \alpha_{i_1-k} e_i + \dots + \\ &+ C_{k-1}^1 A B^{k-2} \sum_{i=2k-2}^n \sum_{i_k-3=2k-2}^{i_k-2} \dots \sum_{i_1=2k-2}^{i_2} \alpha_{i+3-i_{k-3}} \dots \alpha_{i_2+3-i_1} \cdot \alpha_{i_1+5-2k} e_i + \\ &+ B^{k-1} \sum_{i=2k-1}^n \sum_{i_k-2=2k-1}^{i_k-1} \dots \sum_{i_1=2k-1}^{i_2} \alpha_{i+3-i_{k-2}} \dots \alpha_{i_2+3-i_1} \alpha_{i_1+4-2k} e_i) = \\ &= (A+B)(A^{k-1}e_k + C_{k-1}^{k-2} A^{k-2} B \alpha_3 e_{k+1} + (C_{k-1}^{k-2} A^{k-2} B \alpha_4 + \\ &+ C_{k-1}^{k-3} A^{k-3} B^2 \sum_{i_1=k+2}^{k+2} \alpha_{k+5-i_1} \cdot \alpha_{i_1+1-k}) e_{k+2} + \dots + (C_{k-1}^{k-2} A^{k-2} B \alpha_{t+2-k} + \\ &+ C_{k-1}^{k-3} A^{k-3} B^2 \sum_{i_1=k+1}^t \alpha_{t+3-i_1} \alpha_{i_1+1-k} + \\ &+ C_{k-1}^{k-4} A^{k-4} B^3 \sum_{i_2=k+3}^t \sum_{i_1=k+3}^{i_2} \alpha_{t+3-i_2} \cdot \alpha_{i_2+3-i_1} \cdot \alpha_{i_1-k} + \dots + \\ &+ C_{k-1}^1 A B^{k-2} \sum_{i_k-3=2k-2}^t \sum_{i_k-4=2k-2}^{i_k-3} \dots \sum_{i_1=2k-2}^{i_2} \alpha_{t+3-i_{k-3}} \cdot \alpha_{i_{k-3}+3-i_{k-4}} \dots \alpha_{i_2+3-i_1} \alpha_{i_1+5-2k} + \\ &+ B^{k-1} \sum_{i_k-2=2k-1}^t \sum_{i_k-3=2k-1}^{i_k-2} \dots \sum_{i_1=2k-1}^{i_2} \alpha_{t+3-i_{k-2}} \cdot \alpha_{i_{k-2}+3-i_{k-3}} \dots \alpha_{i_2+3-i_1} \cdot \alpha_{i_1+4-2k}) e_t + \\ &+ \dots + (C_{k-1}^{k-2} A^{k-2} B \alpha_{n+2-k} + C_{k-1}^{k-3} A^{k-3} B^2 \sum_{i_1=k+1}^n \alpha_{n+3-i_1} \alpha_{i_1+1-k} + \\ &+ C_{k-1}^{k-4} A^{k-4} B^3 \sum_{i_2=k+3}^n \sum_{i_1=k+3}^{i_2} \alpha_{n+3-i_2} \cdot \alpha_{i_2+3-i_1} \cdot \alpha_{i_1-k} + \dots + \\ &+ C_{k-1}^1 A B^{k-2} \sum_{i_k-3=2k-2}^n \sum_{i_k-4=2k-2}^{i_k-3} \dots \sum_{i_1=2k-2}^{i_2} \alpha_{n+3-i_{k-3}} \cdot \alpha_{i_{k-3}+3-i_{k-4}} \dots \alpha_{i_2+3-i_1} \cdot \alpha_{i_1+5-2k} + \\ &+ B^{k-1} \sum_{i_k-2=2k-1}^n \sum_{i_k-3=2k-1}^{i_k-2} \dots \sum_{i_1=2k-1}^{i_2} \alpha_{n+3-i_{k-2}} \cdot \alpha_{i_{k-2}+3-i_{k-3}} \dots \alpha_{i_2+3-i_1} \alpha_{i_1+4-2k}) e_n) = \\ &= (A+B)(A^{k-1}e_k + \sum_{t=k+1}^n (C_{k-1}^{k-2} A^{k-2} B \alpha_{t+2-k} + C_{k-1}^{k-3} A^{k-3} B^2 \sum_{i_1=k+2}^t \alpha_{t+3-i_1} \cdot \alpha_{i_1+1-k} + \\ &+ C_{k-1}^{k-4} A^{k-4} B^3 \sum_{i_2=k+3}^t \sum_{i_1=k+3}^{i_2} \alpha_{t+3-i_2} \cdot \alpha_{i_2+3-i_1} \cdot \alpha_{i_1-k} + \dots + \\ &+ C_{k-1}^1 A B^{k-2} \sum_{i_k-3=2k-2}^t \sum_{i_k-4=2k-2}^{i_k-3} \dots \sum_{i_1=2k-2}^{i_2} \alpha_{t+3-i_{k-3}} \cdot \alpha_{i_{k-3}+3-i_{k-4}} \dots \alpha_{i_2+3-i_1} \cdot \alpha_{i_1+5-2k} + \\ &+ B^{k-1} \sum_{i_k-2=2k-1}^t \sum_{i_k-3=2k-1}^{i_k-2} \dots \sum_{i_1=2k-1}^{i_2} \alpha_{t+3-i_{k-2}} \cdot \alpha_{i_{k-2}+3-i_{k-3}} \dots \alpha_{i_2+3-i_1} \cdot \alpha_{i_1+4-2k}) e_t). \end{aligned}$$

Consider the products in $L'(\alpha'_3, \alpha'_4, \dots, \alpha'_n, \theta')$:

$$[e'_0, e'_1] = \sum_{k=3}^{n-1} \alpha'_k e'_k + \theta' e'_n, \quad [e'_1, e'_1] = \sum_{k=3}^n \alpha'_k e'_k.$$

Substituting expression e'_k and the lemma 3.3 (for $p=1$), we derive equalities:

$$\begin{aligned} [e'_0, e'_1] &= \sum_{k=3}^{n-1} \alpha'_k (A+B)(A^{k-1}e_k + \sum_{t=k+1}^n (C_{k-1}^{k-2} A^{k-2} B \alpha_{t+2-k} + \\ &+ C_{k-1}^{k-3} A^{k-3} B^2 \sum_{i_1=k+2}^t \alpha_{t+3-i_1} \cdot \alpha_{i_1+1-k} + \\ &+ C_{k-1}^{k-4} A^{k-4} B^3 \sum_{i_2=k+3}^t \sum_{i_1=k+3}^{i_2} \alpha_{t+3-i_2} \cdot \alpha_{i_2+3-i_1} \cdot \alpha_{i_1-k} + \dots + \\ &+ C_{k-1}^1 A B^{k-2} \sum_{i_{k-3}=2k-2}^t \sum_{i_{k-4}=2k-2}^{i_{k-3}} \dots \sum_{i_1=2k-2}^{i_2} \alpha_{t+3-i_{k-3}} \alpha_{i_{k-3}+3-i_{k-4}} \dots \alpha_{i_2+3-i_1} \alpha_{i_1+5-2k} + \\ &+ B^{k-1} \sum_{i_{k-2}=2k-1}^t \sum_{i_{k-3}=2k-1}^{i_{k-2}} \dots \sum_{i_1=2k-1}^{i_2} \alpha_{t+3-i_{k-2}} \alpha_{i_{k-2}+3-i_{k-3}} \dots \alpha_{i_2+3-i_1} \alpha_{i_1+4-2k}) e_t + \\ &+ \theta' A^{n-1} (A+B) e_n = (A+B) \left(\sum_{k=3}^{n-1} A^{k-1} \alpha'_k e_k + \sum_{k=3}^{n-1} \sum_{t=k+1}^n (C_{k-1}^{k-2} A^{k-2} B \alpha_{t+2-k} + \right. \\ &+ C_{k-1}^{k-3} A^{k-3} B^2 \sum_{i_1=k+2}^t \alpha_{t+3-i_1} \alpha_{i_1+1-k} + C_{k-1}^{k-4} A^{k-4} B^3 \sum_{i_2=k+3}^t \sum_{i_1=k+3}^{i_2} \alpha_{t+3-i_2} \alpha_{i_2+3-i_1} \alpha_{i_1-k} + \\ &+ \dots + C_{k-1}^1 A B^{k-2} \sum_{i_{k-3}=2k-2}^t \sum_{i_{k-4}=2k-2}^{i_{k-3}} \dots \sum_{i_1=2k-2}^{i_2} \alpha_{t+3-i_{k-3}} \alpha_{i_{k-3}+3-i_{k-4}} \dots \alpha_{i_2+3-i_1} \alpha_{i_1+5-2k} + \\ &+ B^{k-1} \sum_{i_{k-2}=2k-1}^t \sum_{i_{k-3}=2k-1}^{i_{k-2}} \dots \sum_{i_1=2k-1}^{i_2} \alpha_{t+3-i_{k-2}} \alpha_{i_{k-2}+3-i_{k-3}} \dots \alpha_{i_2+3-i_1} \alpha_{i_1+4-2k}) \alpha'_k e_t + \\ &+ \theta' A^{n-1} e_n \Big) = (A+B) \left(A^2 \alpha'_3 e_3 + \sum_{t=3}^{n-1} A^{t-1} \alpha'_t e_t + \sum_{t=3}^{n-1} \sum_{k=3}^{t-1} (C_{k-1}^{k-2} A^{k-2} B \alpha_{t+2-k} + \right. \\ &+ C_{k-1}^{k-3} A^{k-3} B^2 \sum_{i_1=k+2}^t \alpha_{t+3-i_1} \alpha_{i_1+1-k} + C_{k-1}^{k-4} A^{k-4} B^3 \sum_{i_2=k+3}^t \sum_{i_1=k+3}^{i_2} \alpha_{t+3-i_2} \alpha_{i_2+3-i_1} \alpha_{i_1-k} + \\ &+ \dots + C_{k-1}^1 A B^{k-2} \sum_{i_{k-3}=2k-2}^t \sum_{i_{k-4}=2k-2}^{i_{k-3}} \dots \sum_{i_1=2k-2}^{i_2} \alpha_{t+3-i_{k-3}} \alpha_{i_{k-3}+3-i_{k-4}} \dots \alpha_{i_2+3-i_1} \alpha_{i_1+5-2k} + \\ &+ B^{k-1} \sum_{i_{k-2}=2k-1}^t \sum_{i_{k-3}=2k-1}^{i_{k-2}} \dots \sum_{i_1=2k-1}^{i_2} \alpha_{t+3-i_{k-2}} \alpha_{i_{k-2}+3-i_{k-3}} \dots \alpha_{i_2+3-i_1} \alpha_{i_1+4-2k}) \alpha'_k e_t + \\ &+ (A+B) (\theta' A^{n-1} + \sum_{k=3}^{n-1} (C_{k-1}^{k-2} A^{k-2} B \alpha_{n+2-k} + C_{k-1}^{k-3} A^{k-3} B^2 \sum_{i_1=k+2}^n \alpha_{n+3-i_1} \alpha_{i_1+1-k} + \\ &+ C_{k-1}^{k-4} A^{k-4} B^3 \sum_{i_2=k+3}^n \sum_{i_1=k+3}^{i_2} \alpha_{n+3-i_2} \alpha_{i_2+3-i_1} \alpha_{i_1-k} + \dots + \\ &+ C_{k-1}^1 A B^{k-2} \sum_{i_{k-3}=2k-2}^n \sum_{i_{k-4}=2k-2}^{i_{k-3}} \dots \sum_{i_1=2k-2}^{i_2} \alpha_{n+3-i_{k-3}} \alpha_{i_{k-3}+3-i_{k-4}} \dots \alpha_{i_2+3-i_1} \alpha_{i_1+5-2k} + \\ &+ B^{k-1} \sum_{i_{k-2}=2k-1}^n \sum_{i_{k-3}=2k-1}^{i_{k-2}} \dots \sum_{i_1=2k-1}^{i_2} \alpha_{n+3-i_{k-2}} \alpha_{i_{k-2}+3-i_{k-3}} \dots \alpha_{i_2+3-i_1} \alpha_{i_1+4-2k}) \alpha'_k e_n \Big) = \\ &= (A+B) \left(A^2 \alpha'_3 e_3 + \sum_{t=3}^{n-1} (A^{t-1} \alpha'_t e_t + \sum_{k=3}^{t-1} (C_{k-1}^{k-2} A^{k-2} B \alpha_{t+2-k} + \right. \\ &+ C_{k-1}^{k-3} A^{k-3} B^2 \sum_{i_1=k+2}^t \alpha_{t+3-i_1} \alpha_{i_1+1-k} + \\ &+ C_{k-1}^{k-4} A^{k-4} B^3 \sum_{i_2=k+3}^t \sum_{i_1=k+3}^{i_2} \alpha_{t+3-i_2} \alpha_{i_2+3-i_1} \alpha_{i_1-k} + \dots + \end{aligned}$$

$$\begin{aligned}
& + C_{k-1}^1 AB^{k-2} \sum_{i_{k-3}=2k-2}^t \sum_{i_{k-4}=2k-2}^{i_{k-3}} \dots \sum_{i_1=2k-2}^{i_2} \alpha_{t+3-i_{k-3}} \alpha_{i_{k-3}+3-i_{k-4}} \dots \alpha_{i_2+3-i_1} \alpha_{i_1+5-2k} + \\
& + B^{k-1} \sum_{i_{k-2}=2k-1}^t \sum_{i_{k-3}=2k-1}^{i_{k-2}} \dots \sum_{i_1=2k-1}^{i_2} \alpha_{t+3-i_{k-2}} \alpha_{i_{k-2}+3-i_{k-3}} \dots \alpha_{i_2+3-i_1} \alpha_{i_1+4-2k} \alpha'_k e_t + \\
& + (\theta' A^{n-1} + \sum_{k=3}^{n-1} (C_{k-1}^{k-2} A^{k-2} B \alpha_{n+2-k} + C_{k-1}^{k-3} A^{k-3} B^2 \sum_{i_1=k+2}^n \alpha_{n+3-i_1} \alpha_{i_1+1-k} + \\
& + C_{k-1}^{k-4} A^{k-4} B^3 \sum_{i_2=k+3}^n \sum_{i_1=k+3}^{i_2} \alpha_{n+3-i_2} \alpha_{i_2+3-i_1} \alpha_{i_1-k} + \dots + \\
& + C_{k-1}^1 AB^{k-2} \sum_{i_{k-3}=2k-2}^n \sum_{i_{k-4}=2k-2}^{i_{k-3}} \dots \sum_{i_1=2k-2}^{i_2} \alpha_{n+3-i_{k-3}} \alpha_{i_{k-3}+3-i_{k-4}} \dots \alpha_{i_2+3-i_1} \alpha_{i_1+5-2k} + \\
& + B^{k-1} \sum_{i_{k-2}=2k-1}^n \sum_{i_{k-3}=2k-1}^{i_{k-2}} \dots \sum_{i_1=2k-1}^{i_2} \alpha_{n+3-i_{k-2}} \alpha_{i_{k-2}+3-i_{k-3}} \dots \alpha_{i_2+3-i_1} \alpha_{i_1+4-2k} \alpha'_k e_n).
\end{aligned}$$

The expression for $[e'_1, e'_1]$ is easily obtainable. If we use a substitute in the expression $[e'_0, e'_1]$ that we put instead θ' the coefficient α'_n , namely

$$\begin{aligned}
[e'_1, e'_1] &= (A+B)(A^2 \alpha'_3 e_3 + \sum_{t=4}^n (A^{t-1} \alpha'_t + \sum_{k=3}^{t-1} (C_{k-1}^{k-2} A^{k-2} B \alpha_{t+2-k} + \\
& + C_{k-1}^{k-3} A^{k-3} B^2 \sum_{i_1=k+2}^t \alpha_{t+3-i_1} \alpha_{i_1+1-k} + C_{k-1}^{k-4} A^{k-4} B^3 \sum_{i_2=k+3}^t \sum_{i_1=k+3}^{i_2} \alpha_{t+3-i_2} \alpha_{i_2+3-i_1} \alpha_{i_1-k} + \\
& + \dots + C_{k-1}^1 AB^{k-2} \sum_{i_{k-3}=2k-2}^t \sum_{i_{k-4}=2k-2}^{i_{k-3}} \dots \sum_{i_1=2k-2}^{i_2} \alpha_{t+3-i_{k-3}} \alpha_{i_{k-3}+3-i_{k-4}} \dots \alpha_{i_2+3-i_1} \alpha_{i_1+5-2k} + \\
& + B^{k-1} \sum_{i_{k-2}=2k-1}^t \sum_{i_{k-3}=2k-1}^{i_{k-2}} \dots \sum_{i_1=2k-1}^{i_2} \alpha_{t+3-i_{k-2}} \alpha_{i_{k-2}+3-i_{k-3}} \dots \alpha_{i_2+3-i_1} \alpha_{i_1+4-2k} \alpha'_k e_t).
\end{aligned}$$

On the other hand

$$\begin{aligned}
[e'_0, e'_1] &= [Ae_0 + Be_1, (A+B)e_1 + B(\theta - \alpha_n)e_{n-1}] = \\
&= (A+B)^2 \sum_{t=3}^{n-1} \alpha_t e_t + (A+B)(A\theta + B\alpha_n)e_n, \\
[e'_1, e'_1] &= [(A+B)e_1 + B(\theta - \alpha_n)e_{n-1}, (A+B)e_1 + B(\theta - \alpha_n)e_{n-1}] = \\
&= (A+B) \sum_{t=3}^n \alpha_t e_t.
\end{aligned}$$

If we compare the coefficients of the basis elements e_t and note the coefficient $A+B$ is different from zero, we obtain the restrictions that were outlined in the first assertion of the theorem.

By using the corollary 4.3, assertion b) of the theorem is proved in the same manner. $\square$

Acknowledgments. The second named author would like to acknowledge the hospitality to the University of Sevilla (Spain) and the support of the grant INTAS - 04-83-3035.

REFERENCES

1. Ayupov Sh.A., Omirov B.A. *On some classes of nilpotent Leibniz algebras.* Siberian Math. J., 1, v. 42, 2001, p.18-29.
2. Balavoin D. *Deformation des algebre de Leibniz.* C. R. Acad. Sci. Paris, T. 319, 1994, Serie I, p.783 - 788.
3. Casas Miras J.M. *Central extensions of Leibniz algebras.* Extracta Mathematicae, v. 13, n. 3, 1998, p.393-397.
4. Dzhumadil'daev A.S., Davydov A.A. *Factor-complex for Leibniz cohomology.* Comm. in Algebra, 29(9), 2001, p.4197-4210.

5. Frabetti A. *Leibniz homology of dialgebras of matrices*.
J. Pure and Appl. Algebra, 129, 1998, p. 123-141.
6. Gao Y. *Leibniz homology of unitary Lie algebras*.
J. Pure and Appl. Algebra, 140, 1999, p.33-56.
7. Gomez J.R., Jimenez-Merchan A., Khakimjanov Y. *Low-dimensional filiform Lie algebras*.
J. Pure and Appl. Algebra, 130, 1998, p. 133-158.
8. Goze M., Khakimjanov Yu. *Nilpotent Lie algebras*.
Kluwer Academic Publishers, v. 361, 1996, 336 p.
9. Loday J.-L. *Une version non commutative des de Lie: les de Leibniz*.
L'Ens. Math., 39, 1993, p.269 - 293.
10. Loday J.-L., Pirashvili T. *Universal enveloping algebras of Leibniz algebras and (co)homology*.
Math. Ann., 296, 1993, p.139 - 158.
11. Mikhalev A.A., Umirbaev U.U. *Subalgebras of free Leibniz algebras*.
Comm. in Algebra, 26(2), 1998, p.435-446.
12. Pojidaev A.P. *Solvability of finite-dimensional n -ary commutative Leibniz algebras of characteristic 0*.
Preprint of Universidade de Sao Paulo, Brasil, 2002, pp. 19.
13. Vergne M. *Cohomologie des de Lie nilpotentes*.
Application de Lie nilpotentes. Bull. Soc. Math. France, v. 98, 1970, p.81 - 116.

JOSÉ RAMÓN GÓMEZ.

DPTO. MATEMÁTICA APLICADA I, UNIVERSIDAD DE SEVILLA, SEVILLA (SPAIN)

E-mail address: `jrgomez@us.es`

B.A.OMIROV.

INSITUTE OF MATHEMATICS, TASHKENT (UZBEKISTAN)

E-mail address: `omirovb@mail.ru`